

Reasons for choosing darunavir/ritonavir 600/100 mg twice daily vs. 800/100 mg once daily in treatment-naïve patients: 10 years data from the ICONA cohort

A. Tavelli^{*1}, M. Palma², S. Lo Caputo³, G. Madeddu⁴, P. Bonfanti⁵, B. Menzaghi⁶, S. Nozza⁷, A. Antinori⁸, R. Termini², A. d'Arminio Monforte⁹ for Icona Foundation Study Group

1 Icona Foundation, Milan, Italy; 2 Janssen-Cilag SpA, Cologno Monzese, Italy; 3 Policlinico di Bari, Bari, Italy; 4 University of Sassari, Sassari, Italy; 5 ASST di Lecco, A. Manzoni Hospital, Lecco, Italy; 6 ASST della Valle Olona, Busto Arsizio Hospital, Busto Arsizio, Italy; 7 San Raffaele Scientific Institute, Milan, Italy; 8 INMI "L.Spallanzani" IRCCS, Rome, Italy; 9 ASST Santi Paolo e Carlo, University of Milan, Milan, Italy; *alessandro.tavelli@fondazioneicona.org

Introduction/Summary

- Darunavir/ritonavir (DRV/r) is indicated as first-line option in persons with low adherence or before the results of the resistance test, due to the high genetic barrier and potency.
- DRV/r 600/100 mg twice daily (BID), is licensed for ART-experienced patients, while for ART-naïve patients the recommended dose is 800/100 mg once daily (QD).
- However, in clinical practice, a non negligible proportion of ART-naïve subjects are given DRV /r BID.

Aim

- The aim of this analysis is to identify patterns of use of DRV/r BID, by analyzing predictors of DRV BID vs DRV QD start as first-line treatment

Study Design and Methods

- All patients from the Icona cohort who started a DRV/r-based cART regimen from ART-Naïve in 2008-2017 were included in this analysis.
- A cross-sectional analysis was performed comparing demographics, clinical and lifestyle factors at the time of DRV/r start, comparing BID vs. QD frequency by chi-square and Wilcoxon test as appropriate.
- Univariable and multivariable logistic regression were performed to identify predictors of DRV/r BID initiation.

Results

- A total of 1503 ART-naïve subjects starting their first cART with DRV/r were included: 1297 subjects started DRV/r QD (86.3%) and 206 DRV/r BID (13.7%).

Patients' characteristics

- 82% male, median age (IQR) 40 years (32-48), 80% Italian, 19% with diagnosis of AIDS CD4 count 256 cells/mm³ (85-412), HIV-RNA 5.1 log₁₀ copies/mL (4.5-5.6), other participants' characteristics at the time of starting DRV/r are shown in Table 1.


Table 1 – Main characteristics of the study population at DRV/r initiation

	DRV/r QD (N= 1,297)	DRV/r BID (N=206)	p-value	Total (N=1,503)
Gender, Male, n(%)	1060 (81.7%)	173 (84.0%)	0.434	1233 (82.0%)
Mode of HIV Transmission, n(%)			0.682	
MSM	607 (46.8%)	89 (43.2%)		696 (46.3%)
IDU	79 (6.1%)	11 (5.3%)		90 (6.0%)
Heterosexual	510 (39.2%)	87 (42.2%)		597 (39.7%)
Other/Unknown	101 (7.8%)	19 (9.2%)		120 (8.0%)
Italian, n(%)	1032 (79.6%)	165 (80.1%)	0.041	1197 (79.6%)
Age, years, median(IQR)	40 (32-48)	43 (35-50)	0.005	40 (32-48)
Months from HIV diagnosis, median(IQR)	1.5 (0.6-12.2)	1.4 (0.5-7.9)	0.175	1.4 (0.6-11.7)
Acute HIV Infection, n(%)	31 (2.4%)	5 (2.4%)	0.974	36 (2.4%)
Site Geographical Position, n(%)			0.001	
North	645 (49.7%)	83 (40.3%)		728 (48.4%)
Center	507 (39.1%)	83 (40.3%)		590 (39.2%)
South	145 (11.2%)	40 (19.4%)		185 (12.3%)
CD4 count [cells/mm ³], median(IQR)	260 (88-415)	215 (60-378)	0.067	256 (85-412)
CD4 count ≤200 cells/mm ³ , n(%)	520 (40.1%)	98 (47.6%)	0.043	618 (41.1%)
CD8 count [cells/mm ³], median(IQR)	838 (534-1221)	745 (469-1216)	0.232	824 (523-1221)
CD4/CD8 ratio, median (IQR)	0.28 (0.14-0.46)	0.27 (0.14-0.44)	0.750	0.28 (0.14-0.46)
HIV-RNA [log ₁₀ cps/mL], median(IQR)	5.11 (4.53-5.62)	5.30 (4.58-5.79)	0.020	5.12 (4.53-5.65)
HIV-RNA >500.000 cps/mL, n (%)	288 (22.2%)	63 (30.6%)	0.008	351 (23.3%)
HCV co-infection, n (%)			0.821	
Negative	1051 (81.0%)	169 (82.0%)		1220 (81.2%)
Positive	128 (9.9%)	21 (10.2%)		149 (9.9%)
Not tested	118 (9.10%)	16 (7.8%)		134 (8.9%)
HBV co-infection, n (%)			0.193	
Negative	1072 (82.6%)	170 (82.5%)		1242 (82.6%)
Positive	50 (3.9%)	16 (6.3%)		63 (4.2%)
Not tested	175 (13.5%)	23 (11.2%)		198 (13.2%)
AIDS diagnosis, n (%)	255 (19.7%)	32 (15.5%)	0.162	287 (19.1%)
CVD diagnosis, n (%)	16 (1.2%)	1 (0.5%)	0.346	17 (1.1%)
Diabetes diagnosis, n (%)	27 (2.0%)	6 (2.9%)	0.450	33 (2.2%)
NADMs, n(%)	28 (2.1%)	2 (1.0%)	0.257	30 (2.0%)
Tot. Cholesterol [mg/dL], median(IQR)	153 (130-180)	152 (123-190)	0.905	153 (129-181)
HDL [mg/dL], median (IQR)	37 (29-46)	34 (27-42)	0.018	37 (29-45)
Glucose [mg/dL], median (IQR)	85 (79-94)	85 (80-95)	0.669	85 (79-94)

ARV regimens

- 80.4% started DRV/r in a triple therapy, 5.6% in dual regimen (81% with RAL) and 14.0% in combination with >3 antiretroviral drugs (69% with RAL and 13% with MRV), with a higher proportion of patients on >3 drugs in the DRV/r BID group (22.3% vs 12.6%, p=0.002). 80.0% starting with TDF/FTC as NRTI backbone, 10.8% with ABC/3TC and 1.1% with AZT/3TC
- There was a peak in the use of DRV/r BID in 2012-2013 (p<0.001) [Figure]

Figure – Proportion of DRV/r use by frequency (QD vs. BID) according to calendar period



Predictors of DRV/r BID start

- Fitting a logistic regression model, after controlling for potential confounders, older patients, subjects with CD4<200/mm³, patients given regimens >3 ARV drugs, patients from centers of southern Italy and patients treated in 2012-2013 showed a significantly higher probability of starting DRV/r BID. A decreased probability of DRV/r BID start was observed in patients with AIDS diagnosis [Table 2]. There was evidence that the use of regimens with >3 ARV was greater in people with HIV RNA >500.000 copies/mL than in those with HIV RNA ≤ 500.000 copies/mL (OR=5.9; p<0.001)

Table 2 – Crude odds ratio (OR) and adjusted odds ratio (AOR) of starting DRV/r BID

	OR	95%CI	p-value	AOR	95%CI	p-value
Gender, Male	1.17	0.78-1.74	0.434	1.23	0.81-1.87	0.329
Mode of HIV Transmission						
Heterosexual	1.00					
IDU	0.82	0.42-1.60	0.553			
MSM	0.86	0.62-1.18	0.351			
Other/Unknown	1.10	0.64-1.89	0.723			
Nationality, Italian	1.06	0.73-1.53	0.761			
Age, per 10 years older	1.20	1.06-1.37	0.004	1.20	1.05-1.37	0.007
N. ARV Drugs Started						
≤3 Drugs	1.00					
>3 Drugs	1.98	1.38-2.86	<.001	2.14	1.42-3.25	<.001
Calendar Period of Start						
2016-2017	1.00					
2014-2015	1.05	0.62-1.79	0.849	1.23	0.72-2.13	0.443
2012-2013	2.01	1.21-3.33	0.006	2.47	1.47-4.17	0.001
≤2011	0.59	0.32-1.08	0.089	0.84	0.44-1.59	0.595
Site Geographical Position						
Northern Italy	1.00					
Central Italy	1.27	0.92-1.76	0.147	1.27	0.91-1.78	0.162
Southern Italy	2.14	1.41-3.26	0.000	2.64	1.68-4.13	<.001
CD4 count, cells/mm ³						
>200	1.00					
≤200	1.35	1.00-1.82	0.043	1.45	1.04-2.03	0.029
HIV-RNA copies/mL, per 1 log ₁₀ increase	1.22	1.03-1.44	0.019	1.05	0.88-1.26	0.546
Hepatitis Co-Infection	0.77	0.40-1.51	0.458			
Acute HIV Infection	1.01	0.39-2.64	0.974			
AIDS diagnosis	0.75	0.50-1.12	0.163	0.61	0.39-0.95	0.030

Conclusion

- Italian clinicians allocate DRV/r BID as first line regimen in several scenarios: elderly subjects, severely immunodepressed, highly viremic patients, more frequently in combination with more than 3 drugs and in patients treated in 2012-2013.
- Subjects with AIDS are mostly given DRV/r QD probably due to the concomitant medications for opportunistic infections.

Funding

- This analysis is supported by Janssen Italy.
- ICONA Foundation is supported by unrestricted grants from BMS, Gilead, Janssen, MSD and ViiV Healthcare

Icona Foundation Study Group